

Food Calorimetry Lab

Introduction


When a substance is heated, the motion of its individual particles increases, resulting in an increase in temperature. The more heat that is added per gram of substance, the greater the temperature change. The relationship between the heat added, the mass of a substance, and the temperature change it undergoes is known as *specific heat*.

$$\text{Specific Heat} = \frac{\text{Energy in joules}}{\text{Mass in grams} \times \text{Temperature change in Celsius or Kelvin}}$$

Specific heat is defined as the amount of energy necessary to produce a temperature change of 1°C per gram of substance. The specific heats of different substances vary, and therefore this quantity may be useful in identifying an unknown.

The measurement of heat changes is called *calorimetry*. A calorimeter is a piece of equipment designed to measure the energy released or absorbed during a chemical reaction or phase change. In this lab, calorimetry will be used to determine the amount of heat produced during the combustion of various food samples. A piece of food will be burned and the released energy will be used to heat a known quantity of water. The temperature change of water will then be used to determine the amount of energy in the food.

The diagram below shows the basic setup of the food calorimeter.



It will be assumed that the entire amount of energy released by the burning food is gained by the water. This is based on the Law of Conservation of Energy, which states that energy is neither created nor destroyed. We will assume no heat loss to the calorimeter or the environment outside the calorimeter.

Energy released by burning food = Energy gained by water

The formula below is used to calculate the amount of energy absorbed/released during calorimetry.

$$q = mc\Delta T$$

where **q** = heat (in joules); **m** = mass (in grams); **c** = specific heat (in joules/grams • °C); **ΔT** = change in temperature (i.e. final temp – initial temp) (in °C or K)

