

Properties of Solids

Name: \_\_\_\_\_ Date: \_\_\_\_\_

Model 1 – Types of Solids













Ionic Solids	 <p data-bbox="293 548 581 583">Iron(II) Sulfide (FeS)</p>	 <p data-bbox="699 548 1049 583">Calcium Bromide (CaBr<sub>2</sub>)</p>	 <p data-bbox="1143 548 1487 583">Sodium Chloride (NaCl)</p>
Molecular Solids	 <p data-bbox="337 919 537 955">Nitrogen (N<sub>2</sub>)</p>	 <p data-bbox="805 919 943 955">Ice (H<sub>2</sub>O)</p>	 <p data-bbox="1214 919 1414 955">Dry Ice (CO<sub>2</sub>)</p>
Network Covalent Solids	 <p data-bbox="305 1283 570 1318">Diamond (Carbon)</p>	 <p data-bbox="792 1283 959 1318">Silica (SiO<sub>2</sub>)</p>	 <p data-bbox="1235 1283 1393 1318">Silicon (Si)</p>
Metallic Solids	 <p data-bbox="342 1646 532 1682">Platinum (Pt)</p>	 <p data-bbox="748 1646 1008 1682">Brass (Cu and Zn)</p>	 <p data-bbox="1235 1646 1393 1682">Nickel (Ni)</p>

Table 1

	<b>A</b>	<b>B</b>	<b>C</b>	<b>D</b>	<b>E</b>	<b>F</b>
	<b>All atoms are nonmetals</b>	<b>All atoms are metals</b>	<b>Atoms are metals and nonmetals</b>	<b>Molecular structure</b>	<b>Formula units</b>	<b>Attractive forces</b>
<b>Ionic Solids</b>						
<b>Molecular Solids</b>						
<b>Network Covalent Solids</b>						
<b>Metallic Solids</b>						

Table 2

	<b>Bonds/forces broken upon melting</b>	<b>Individual particles in the liquid</b>
<b>Ionic</b>		
<b>Molecular</b>		
<b>Network covalent</b>		
<b>Metallic</b>		

## Model 2 – Melting Points and Enthalpies of Fusion

Type of Solid	Substance	Chemical Formula	Melting Point (°C)	Enthalpy of Fusion (kJ/mole)
Ionic	Iron(II) sulfide	FeS	1195	51.0
	Calcium bromide	CaBr <sub>2</sub>	730	17.5
	Sodium chloride	NaCl	804	30.3
Molecular	Nitrogen	N <sub>2</sub>	-210	0.72
	Water	H <sub>2</sub> O	0.0	6.02
	Carbon dioxide	CO <sub>2</sub>	-78	8.10
Network covalent	Diamond	C	3550	117.0
	Silica	SiO <sub>2</sub>	1650	12.5
	Silicon	Si	1687	50.0
Metallic	Platinum	Pt	1770	24.0
	Brass	Cu and Zn	~930	Varies
	Nickel	Ni	1453	71.0

1. If both molecular and network covalent solids contain covalent bonds, then why are the melting points of molecular solids so much lower than those of network covalent solids?

### Model 3 – Solubility and Conductivity of Solids

Substance	Chemical Formula	Solubility in Water	Conductive as a Solid	Conductive as a Liquid	Conductive in Aqueous Solution
Iron(II) sulfide	FeS	Insoluble	No	Yes	N/A
Calcium bromide	CaBr <sub>2</sub>	Very soluble	No	Yes	Yes
Sodium chloride	NaCl	Very soluble	No	Yes	Yes
Nitrogen	N <sub>2</sub>	Slightly soluble	No	No	No
Water	H <sub>2</sub> O	N/A	No	Slightly	N/A
Carbon dioxide	CO <sub>2</sub>	Slightly soluble	No	No	No
Diamond	C	Insoluble	No	No	N/A
Silica	SiO <sub>2</sub>	Insoluble	No	No	N/A
Platinum	Pt	Insoluble	Yes	Yes	N/A
Brass	Cu and Zn	Insoluble	Yes	Yes	N/A
Nickel	Ni	Insoluble	Yes	Yes	N/A

N/A = not applicable

2. In general, which type of solid is very soluble in water? Explain your reasoning.

3. Why are ionic compounds conductive as liquids, but not as solids?

4. Why are metals conductive as solids?

Metallic Sea of Electrons

